

# Let it Glow HOLIDAY PROMOTION

## SPARKLE. SHINE. SAVE.

This holiday season, get a naturally youthful look for less with *Dysport®* (abobotulinumtoxinA) for Injection® and the *Restylane®* family of fillers.† Smooth frown lines between the brows with *Dysport*, and achieve youthfullooking results with *Restylane*-and save!

The most common side effects for *Dysport* are nose and throat irritation, headache, injection site pain, injection site skin reaction, upper respiratory tract infection, eyelid swelling, eyelid drooping, sinus inflammation and nausea.

\*Please see full Important Safety Information, including Distant Spread of Toxin Effect Boxed Warning, on the back of this piece.

<sup>†</sup>Offer terms and conditions apply. See details on the back of this piece.

Actual patient, Individual results may vary.

#### Let It Glow Offer Consumer Terms & Conditions

This offer is limited to adult patients who receive a *Dysport\** treatment and a *Restylane\**, *Restylane-L\**, *Restylane\*Silk*, *Restylane\*Lyft*, *Restylane\* Refyne*, or *Restylane\*Defyne* (a "Qualifying HA Filler") treatment in a single office visit in accordance with the products' FDA-approved indications and labeling. All 0.5 ml syringes of Qualifying HA filler are excluded from this offer. You must be a member of the ASPIRE Galderma Rewards program. This is a limited quantity offer subject to availability from Galderma and your participating healthcare provider. If you have questions about this offer, please call 1-844-5ASPIRE.

To participate in this offer, you must receive at least a qualifying treatment of *Dysport* and a qualifying treatment with one (1) syringe of Qualifying HA Filler in a single office visit on or after October 7, 2019, and your healthcare provider must submit a valid treatment certificate coupon code to the ASPIRE Galderma Rewards program on your behalf at the time of treatment. Your healthcare provider may submit: a \$100 or \$125 coupon code if you receive a qualifying treatment of *Dysport* and a qualifying treatment with one (1) syringe of Qualifying HA Filler, or a \$150 or \$175 coupon code if you receive a qualifying treatment of *Dysport* and a qualifying treatment with two (2) syringes of Qualifying HA Filler. Whether your healthcare provider may submit a \$100 or \$125 code, or a \$150 or \$175 code, is determined by your healthcare provider's prior purchases of *Dysport* and Qualifying HA Filler from Galderma. Coupon codes shall expire sixty (60) days from the date issued to the healthcare provider.

You are eligible for this offer only if you pay for your entire treatment yourself and if no part of your treatment is covered by insurance or any other third party payor. This offer excludes any treatment using a Galderma product that is reimbursed by Medicaid, Medicare, or any other federal or state benefit programs, including state medical assistance programs. You are not eligible for this offer if your private insurance, HMO, or any other health benefit program pays for all or part of their treatment. If any form of reimbursement is sought from a third-party, you may be required to disclose the value of the coupon code to that party. This offer is not available to patients who are Galderma employees or their spouses.

This offer is non-transferrable and available and valid only from participating healthcare providers. All treatment certificate coupon codes must be submitted by the treating provider via <a href="www.aspirehcp.com">www.aspirehcp.com</a>. Galderma will reimburse the provider for the face value of used coupon codes, provided that the provider and patient have complied with these terms and conditions. Individual treatment certificate coupon codes are valid for one-time use. This offer is limited to one (1) treatment certificate coupon code per patient. Promotion treatment certificate coupon codes cannot be combined with any other offer or promotion except treatment certificates created by the patient through the redemption of ASPIRE Galderma Rewards program points. Treatment certificate coupon codes have no cash value without simultaneous purchase of qualifying <a href="https://doi.org/10.1081/jusport">Dysport</a> and a Qualifying HA Filler treatments, and are void if sold, transferred or exchanged. Offer valid only in the U.S., excluding territories, and void where prohibited, taxed or otherwise restricted by law.

By participating in this offer, you agree to all of the terms and conditions of this offer. Galderma reserves the right to cancel or modify this offer without notice. Galderma assumes no responsibility for lost, late, incomplete, or incorrect submissions that fail to be properly entered and submitted to <a href="https://www.aspirehcp.com">www.aspirehcp.com</a> for any reason. Treatment certificate coupon codes will be processed and reimbursed in U.S. dollars only.

#### Dysport

Dysport is a prescription injection for temporary improvement in the look of moderate to severe frown lines between the eyebrows (glabellar lines) in adults less than 65 years of age.

#### Important Safety Information

What is the most important information you should know about *Dysport*? Spread of Toxin Effects: In some cases, the effects of *Dysport* and all botulinum toxin products may affect areas of the body away from the injection site. Symptoms can happen hours to weeks after injection and may include swallowing and breathing problems, loss of strength and muscle weakness all over the body, double vision, blurred vision and drooping eyelids, hoarseness or change or loss of voice, trouble saying words clearly, or loss of bladder control. Swallowing and breathing problems can be life threatening and there have been reports of death. You are at the highest risk if these problems are pre-existing before injection.

These effects could make it unsafe for you to drive a car, operate machinery, or do other dangerous activities.

Do not have Dysport treatment if you: are allergic to Dysport or any of its ingredients (see the end of the Medication Guide for a list of ingredients), are allergic to cow's milk protein, had an allergic reaction to any other botulinum toxin product, such as Myobloc\*, Botox\*, or Xeomin\*, have a skin infection at the planned injection site, under 18 years of age, or are pregnant or breastfeeding.

The dose of *Dysport* is not the same as the dose of any other botulinum toxin product and cannot be compared to the dose of any other product you may have used.

Tell your doctor about any swallowing or breathing difficulties and all your muscle or nerve conditions such as amyotrophic lateral sclerosis [ALS or Lou Gehrig's disease], myasthenia gravis, or Lambert-Eaton syndrome, which may increase the risk of serious side effects including difficulty swallowing and difficulty breathing.

**Tell your doctor about all of your medical conditions**, including if you have surgical changes to your face, very weak muscles in the treatment area, any abnormal facial change, injection site inflammation, droopy eyelids or sagging eyelid folds, deep facial scars, thick oily skin, wrinkles that can't be smoothed by spreading them apart, or if you are pregnant or breastfeeding or planning to become pregnant or breastfeed.

Tell your doctor about all the medicines you take, including prescription and nonprescription medicines, vitamins and herbal and other natural products. Using *Dysport* with certain other medicines may cause serious side effects. Do not start any new medicines while taking *Dysport* without talking to your doctor first.

Especially tell your doctor if you: have received any other botulinum toxin product, such as Myobloc® (rimabotulinumtoxinB), Botox® (onabotulinumtoxinA), or Xeomin® (incobotulinumtoxinA), in the last four months or any in the past (be sure your doctor knows exactly which product you received, have recently received an antibiotic by injection, take muscle relaxants, take an allergy or cold medicine, or take a sleep medicine.

#### **Common Side Effects**

The most common side effects are nose and throat irritation, headache, injection site pain, injection site skin reaction, upper respiratory tract infection, eyelid swelling, eyelid drooping, sinus inflammation, and nausea.

Ask your doctor if Dysport is right for you.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit <a href="www.fda.gov/medwatch">www.fda.gov/medwatch</a> or call 1-800-FDA-1088.

Please use  ${\it Dysport}$  Full Prescribing Information including Medication Guide at  ${\it DysportUSA.com}$ .

#### Restylane

#### Important Safety Information

The Restylane family of products includes Restylane\*, Restylane-L\*, Restylane\* Lyft with Lidocaine, Restylane\* Silk, Restylane\* Refyne, and Restylane\* Defyne.

#### APPROVED USES

Restylane\* and Restylane-L\* are for mid-to-deep injection into the facial tissue for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds. Restylane\* and Restylane-L\* are also indicated for injection into the lips in patients over the age of 21.

Restylane\* Lyft with Lidocaine is for deep implantation into the facial tissue for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds and for cheek augmentation and for the correction of agerelated midface contour deficiencies in patients over the age of 21. Restylane\* Lyft with Lidocaine is also indicated for injection into the subcutaneous plane in the dorsal hand to correct volume deficit in patients over the age of 21.

Restylane\* Silk is for lip augmentation and for correction of perioral wrinkles in patients over the age of 21.

Restylane\* Refyne is for mid-to-deep injection into the facial tissue for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds, in patients over the age of 21.

Restylane® Defyne is for mid-to-deep injection into the facial tissue for the correction of moderate to severe deep facial wrinkles and folds, such as nasolabial folds, in patients over the age of 21.

### Are there any reasons why I should not use products within the Restylane\* family? (Contraindications)

To ensure a safe procedure, your doctor will talk to you about your medical history to determine if you are an appropriate candidate for treatment. You should not use products within the Restylane family if:

- You have severe allergies with a history of severe reactions (anaphylaxis)
- You are allergic to lidocaine or to any of the gram-positive bacterial proteins used to make hyaluronic acid
- · You are prone to bleeding or have been diagnosed with a bleeding disorder

#### Are there other precautions that I should discuss with my doctor?

- Tell your doctor if you are breastfeeding, pregnant, or trying to become pregnant. The safety of these products for use during pregnancy, or in women who are breastfeeding, has not been studied
- Restylane, Restylane-L, Restylane\* Lyft with Lidocaine, Restylane Refyne
  and Restylane Defyne are intended to treat facial wrinkles and folds, such
  as nasolabial folds. Restylane and Restylane-L are also intended for lip
  enhancement. Restylane\* Lyft with Lidocaine is also intended for injection in
  the dorsal hand to correct volume loss. Treatments in other areas of the face
  or body have not been evaluated in clinical studies.
- The safety and effectiveness of Restylane\* Silk for areas other than the lips and perioral area have not been evaluated in clinical studies.
- Tell your doctor if you have any history of scarring, particularly thick and stiff scars, or any pigmentation (skin color) disorders. These side effects can occur with hyaluronic acid fillers in general.
- Tell your doctor if you are planning other laser treatments or a chemical peel, as there is a possible risk of inflammation at the treatment site if these procedures are performed after treatment.
- Patients who experience skin injury near the site of injection with these products may be at a higher risk for side effects
- Tell your doctor if you are on any medications to decrease your body's immune response (immunosuppressive therapy). Using these medications may increase your risk of bruising or bleeding at the gel injection site.
- Tell your doctor if you are using any "blood thinners" such as aspirin, warfarin, or any other medications that affect bleeding. Using these medications may increase your risk of bruising or bleeding at the gel injection site.
- The use of these products on gel injection sites with skin sores, pimples, rashes, hives, cysts, or infections should be postponed until healing is complete. Use of product in these areas could delay healing or make your skin problems worse.
- Tell your doctor if you have diseases, injuries, or disabilities of the hand.

#### What are the possible side effects?

The most commonly observed side effects are swelling, redness, pain, bruising, headache, tenderness, lump formation, itching at the injection site, and impaired hand function. These are typically mild in severity and typically resolve in less than 7 days in nasolabial folds and less than 14 days in lips. Serious but rare side effects include delayed onset infections, recurrence of herpetic eruptions, and superficial necrosis at the injection site.

One of the risks with using this product is unintentional injection into a blood vessel. The chances of this happening are very small, but if it does happen, the complications can be serious, and may be permanent. These complications, which have been reported for facial injections, can include vision abnormalities, blindness, stroke, temporary scabs, or permanent scarring of the skin.

As with all skin injection procedures, there is a risk of infection.

To report a side effect with any of the Restylane products, please call Galderma Laboratories, L.P at 1-855-425-8722.

The Restylane family of products is available only through a licensed practitioner. Complete Instructions for Use are available at www.RestylaneUSA.com.

© 2019 Galderma Laboratories, L.P. The Dysport trademark is used under license

All trademarks are the property of their respective owners.

USMP/DYS/0230/0919b

